

IN THE CLAIMS

Please amend claims 116 and 164, as shown below. Please cancel claims 132-137, 142-145, 152-159, 161-163, 167, and 175-177 without prejudice. The following listing of claims replaces all prior listings.

1-115. (Canceled).

116. (Currently amended) A method for enhancing the delivery of a bioactive agent from the vasculature to a selected tissue in a patient, comprising:

(i) administering said bioactive agent to said patient;

(ii) administering a vesicle composition to said patient, by intravascular infusion, wherein said vesicle composition comprises, in an aqueous carrier, vesicles comprising lipids, proteins, or polymers and a gas or gaseous precursor; and

(iii) cavitating and/or rupturing said vesicles by applying to the patient ultrasonic energy having a frequency between about 750 kHz and 3 MHz, to the patient in an amount sufficient to produce cavitation or rupture of said vesicles, and sufficient to increase thereby enhance delivery of said bioactive agent from the vasculature through the vessel wall and into said selected tissue, ~~wherein said bioactive agent is delivered into said selected tissue, wherein said ultrasound energy has a frequency between about 750 kHz and 3 MHz.~~

117. (Previously presented) A method according to Claim 116 wherein said bioactive agent is administered to said patient at a rate which comprises continuous infusion.

118. (Previously presented) A method according to Claim 116 wherein said bioactive agent and said vesicle composition are administered to said patient substantially simultaneously.

119. (Previously presented) A method according to Claim 116 further comprising imaging said patient using diagnostic ultrasound imaging.

120. (Previously presented) A method according to Claim 116 wherein said vesicles comprise lipids.

121. (Previously presented) A method according to Claim 120 wherein said vesicle composition comprises vesicles selected from the group consisting of micelles and liposomes.

122. (Previously presented) A method according to Claim 120 wherein said lipids comprise phospholipids.

123. (Previously presented) A method according to Claim 122 wherein said phospholipids are selected from the group consisting of phosphatidylcholine, phosphatidylethanolamine and phosphatidic acid.

124. (Previously presented) A method according to Claim 123 wherein said phosphatidylcholine is selected from the group consisting of dioleoylphosphatidylcholine, dimyristoylphosphatidylcholine, dipalmitoylphosphatidylcholine and distearoylphosphatidylcholine.

125. (Previously presented) A method according to Claim 124 wherein said phosphatidylcholine comprises dipalmitoylphosphatidylcholine.

126. (Previously presented) A method according to Claim 123 wherein said phosphatidylethanolamine is selected from the group consisting of dipalmitoylphosphatidylethanolamine, dioleoylphosphatidylethanolamine, N-succinyldioleoylphosphatidylethanolamine and 1-hexadecyl-2-palmitoylglycerophosphoethanolamine.

127. (Previously presented) A method according to Claim 126 wherein said phosphatidylethanol-amine comprises dipalmitoylphosphatidylethanolamine.

128. (Previously presented) A method according to Claim 123 wherein said phosphatidic acid comprises dipalmitoylphosphatidic acid.

129. (Previously presented) A method according to Claim 120 wherein said lipid further comprises a polymer.

130. (Previously presented) A method according to Claim 129 wherein said polymer comprises a hydrophilic polymer.

131. (Previously presented) A method according to Claim 130 wherein said hydrophilic polymer comprises polyethylene glycol.

132-137. (Canceled).

138. (Previously presented) A method according to Claim 116 wherein said gas comprises a fluorinated gas.

139. (Previously presented) A method according to Claim 138 wherein said fluorinated gas is selected from the group consisting of a perfluorocarbon and sulfur hexafluoride.

140. (Previously presented) A method according to Claim 139 wherein said fluorinated gas comprises a perfluorocarbon.

141. (Previously presented) A method according to Claim 140 wherein said perfluorocarbon gas is selected from the group consisting of perfluoromethane, perfluoroethane, perfluoropropane, perfluorobutane and perfluorocyclobutane.

142-145. (Canceled).

146. (Previously presented) A method according to Claim 116 wherein said vesicle composition is administered to the patient at a rate of from about 1×10^6 to less than about 8×10^6 vesicles/Kg-sec.

147. (Previously presented) A method according to Claim 146 wherein said vesicle composition is administered at a rate of from about 1×10^6 to about 7×10^6 vesicles/Kg-sec.

148. (Previously presented) A method according to Claim 147 wherein said vesicle composition is administered at a rate of from about 1.5×10^6 to about 6×10^6 vesicles/Kg-sec.

149. (Previously presented) A method according to Claim 148 wherein said vesicle composition is administered at a rate of from about 2×10^6 to about 5.5×10^6 vesicles/Kg-sec.

150. (Previously presented) A method according to Claim 149 wherein said vesicle composition is administered at a rate of from about 2.5×10^6 to about 5×10^6 vesicles/Kg-sec.

151. (Previously presented) A method according to Claim 150 wherein said vesicle composition is administered at a rate of from about 3×10^6 to about 4.5×10^6 vesicles/Kg-sec.

152-159.(Canceled).

160. (Previously presented) A method according to Claim 116 wherein said bioactive agent is selected from the group consisting of a diagnostic agent, genetic material, a peptide, a beta-agonist, an anti-asthmatic, a steroid, a cholinergic agent, an anti-cholinergic agent, a 5-lipoxygenase inhibitor, a leukotriene inhibitor, an anti-neoplastic agent, an antibiotic, an anti-tumor drug, a radiation sensitizer, a thrombolytic agent, an

anti-histamine, an anti-coagulant, an anti-inflammatory, a hormone, a growth factor, an angiogenic factor and a mitotic inhibitor.

161. (Canceled).

164. (Currently amended) A method for enhancing the delivery of a bioactive agent from the vasculature to a selected tissue in a patient, said method comprising:

(i) administering said bioactive agent to said patient;

(ii) administering an acoustically active composition to said patient, by intravascular infusion; and

(iii) activating said acoustically active composition by applying to the patient ultrasonic energy having a frequency between about 750 kHz and 3 MHz, to the patient in an amount sufficient to activate said acoustically active composition, and sufficient to increase thereby enhance delivery of said bioactive agent from the vasculature into said selected tissue, ~~wherein said bioactive agent is delivered into said selected tissue and said ultrasound energy has a frequency of from about 750 kHz to 3 MHz.~~

165. (Previously presented) A method according to Claim 164 wherein said bioactive agent is administered to said patient at a rate which comprises continuous infusion.

166. (Previously presented) A method according to Claim 164 wherein said bioactive agent and said acoustically active composition are administered to said patient substantially simultaneously.

167. (Canceled).

168. (Previously presented) A method according to Claim 164 wherein said tissue comprises an area of reduced blood perfusion.

169. (Previously presented) A method according to Claim 168 wherein said area of reduced blood perfusion comprises ischemic tissue.

170. (Previously presented) A method according to Claim 164 wherein said tissue comprises myocardium.

171. (Previously presented) A method according to Claim 164 wherein said tissue comprises glandular tissue.

172. (Previously presented) A method according to Claim 171 wherein said glandular tissue comprises the prostate gland.

173. (Previously presented) A method according to Claim 164 further comprising imaging said tissue using diagnostic ultrasound imaging.

174. (Previously presented) A method according to Claim 164 wherein said bioactive agent comprises an agent selected from the group consisting of a diagnostic agent, genetic material, a peptide, a beta-agonist, an anti-asthmatic, a steroid, a cholinergic agent, an anti-cholinergic agent, a 5-lipoxygenase inhibitor, a leukotriene inhibitor, an anti-neoplastic agent, an antibiotic, an anti-tumor drug, a radiation sensitizer, a thrombolytic agent, an anti-histamine, an anti-coagulant, an anti-inflammatory, a hormone, a growth factor, an angiogenic factor and a mitotic inhibitor.

175-177. (Canceled).

178. (Previously presented) A method according to Claim 174 wherein said acoustically active composition and bioactive agent are administered prior to said application of ultrasound energy.

179. (Previously presented) A method according to Claim 174 wherein said acoustically active composition and bioactive agent are administered at about the same time as said application of ultrasound energy.

180. (Previously presented) A method according to Claim 174 further comprising applying radiation energy to said tissue.

181. (Previously presented) A method according to Claim 179 wherein said acoustically active composition and bioactive agent are administered prior to said application of radiation energy.

182. (Previously presented) A method according to Claim 180 wherein said acoustically active composition and bioactive agent are administered at about the same time as said application of radiation energy.

183. (Previously presented) A method according to Claim 178 wherein said acoustically active composition and bioactive agent are administered from about 1 minute to about 8 hours prior to said application of ultrasound energy.

184. (Previously presented) A method according to Claim 181 wherein said acoustically active composition and bioactive agent are administered from about 1 minute to about 8 hours prior to said application of radiation energy.